



Rare:	Hyperkalaemia, hypomagnesaemia
Psychiatric disorders	

Common:	Insomnia
Uncommon:	Depression, libido decreased, anxiety
Rare:	Confusional state
Nervous system disorders	
Very common:	Headache ²
Common:	Dizziness, paraesthesia, taste disturbance, hyposaesthesia
Uncommon:	Migraine, somnolence, syncope, peripheral neuropathy, memory impairment, sciatica, restless leg syndrome, tremor, cerebral haemorrhage
Rare:	Increased intracranial pressure, convulsions, optic neuritis
Ear and labyrinth disorders	
Uncommon:	Vertigo, tinnitus, hearing loss
Cardiac disorder	
Uncommon:	Palpitations, tachycardia, cardiac failure, congestive3, pulmonary oedema
Rare:	Arrhythmia, atrial fibrillation, cardiac arrest, myocardial infarction, angina pectoris, pericardial effusion
Vascular disorders4	
Common:	Flushing, haemorrhage
Uncommon:	Hypertension, haematoma, subdural haematoma, peripheral coldness, hypotension, R aynaud's phenomenon
Respiratory, thoracic and mediastinal disorders	
Common:	Dyspnoea, epistaxis, cough
Uncommon:	Pleural effusion, pharyngolaryngeal pain, pharyngitis
Rare:	Pleuritic pain, pulmonary fibrosis, pulmonary hypertension, pulmonary haemorrhage
Gastrointestinal disorders	
Very common:	Nausea, diarrhoea, vomiting, dyspepsia, abdominal pain
Common:	Flatulence, abdominal distension, gastro-oesophageal reflux, constipation, dry mouth, gastritis
Rare:	Stomatitis, mouth ulceration, gastrointestinal haemorrhage, eructation, melena, esophagitis, ascites, gastric ulcer, haematemesis, cheilitis, dysphagia, pancreatitis
Hepatobiliary disorders	
Common:	Increased hepatic enzymes
Uncommon:	Hyperbilirubinaemia, hepatitis, jaundice
Rare:	Hepatic failure ⁵ , hepatic necrosis ⁵
Skin and subcutaneous tissue disorders	
Very common:	Periorbital edema, dermatitis/eczema/rash
Uncommon:	Rash pustular, contusion, sweating increased, urticaria, ecchymosis, increased tendency to bruise, hypotrichosis, skin hypopigmentation, dermatitis exfoliative, onychoclasia, folliculitis, petechiae, psoriasis, purpura, skin hyperpigmentation, bullous eruptions
Rare:	Acute febrile neutrophilic dermatosis (Sweet's syndrome), nail discolouration, angioneurotic oedema, rash vesicular, erythema multiforme, leucocytoclastic vasculitis, Stevens-Johnson syndrome, acute generalised exanthematous pustulosis (AGEP)
Musculoskeletal and connective tissue disorders	
Very common:	Muscle spasm and cramps, Musculoskeletal pain including myalgia, arthralgia, bone pain ⁶
Common:	Joint swelling
Uncommon:	Joint and muscle stiffness
Rare:	Muscular weakness, arthritis
Renal and urinary disorders	
Uncommon:	Renal pain, haematuria, renal failure acute, urinary frequency increased
Reproductive system and breast disorders	
Uncommon:	Gynaecomastia, erectile dysfunction, menorrhagia, menstruation irregular, sexual dysfunction, nipple pain, breast enlargement, scrotal oedema
General disorders and administration site conditions	
Very common:	Fluid retention and oedema, fatigue
Investigations	
Very Common:	Weight increased
Common:	Weight decreased
Uncommon:	Blood creatinine increased, blood creatine phosphokinase increased, blood lactate dehydrogenase
Rare:	Blood amylase increased

1.Pneumonia was reported most commonly in patients with transformed CML and in patients with GIST. 2.Headache was the most common in GIST patients. 3. On a patient-year basis, cardiac events including congestive heart failure were more commonly observed in patients with transformed CML than in patients with chronic CML. 4. Flushing was most common in GIST patients and bleeding (hematoma, hemorrhage) was most common in patients with GIST and with transformed CML (CML-AP and CML-BC). 5. Pleural effusion was reported more commonly in patients with GIST and in patients with transformed CML (CML-AP and CML-BC) than in patients with chronic CML. 6. Abdominal pain and gastrointestinal hemorrhage were most commonly observed in GIST patients. 7. Musculoskeletal pain and related events were more commonly observed in patients with CML than in GIST patients. 8. Some fatal cases of hepatic failure and of hepatic necrosis have been reported. The following types of ADRs have been reported from post-marketing experience and from additional clinical studies with Imatinib. They include spontaneous case reports as well as serious ADRs from smaller or ongoing clinical studies and the expanded access programs. Because these ADRs are reported from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to Imatinib exposure.

Table 3 Adverse drug reactions from post-marketing reports

Investigations	
Not known:	Not known: Hepatitis B reactivation
Nervous system disorders	
Uncommon	Cerebral oedema
Eye disorders	
Rare:	Vitreous haemorrhage
Cardiac disorders	
Rare:	Pericarditis, cardiac tamponade

Vascular disorders	
Uncommon:	Thrombosis/embolism
Very rare:	Anaphylactic shock
Respiratory, thoracic and mediastinal disorders	
Uncommon:	Acute respiratory failure ⁷ , interstitial lung disease
Gastrointestinal disorders	
Uncommon:	Ileus/intestinal obstruction, tumour haemorrhage/tumour necrosis, gastrointestinal perforation ⁷
Rare:	Diverticulitis, gastric antral vascular ectasia (GAVE)
Skin and subcutaneous tissue disorders	
Uncommon:	Palmar-plantar erythrodysaesthesia syndrome
Rare:	Lichenoid keratosis, lichen planus, pemphigus
Very rare:	Toxic epidermal necrolysis
Not known:	Drug rash with eosinophilia and systemic symptoms (DRESS), pseudoporphyria
Musculoskeletal and connective tissue disorders	
Verycommon:	Musculoskeletal pain upon treatment discontinuation (including myalgia, pain in extremity, arthralgia, bone pain, spinal pain)
Uncommon:	Osteonecrosis
Rare:	Rhabdomyolysis/myopathy
Not known:	Growth retardation in children
Reproductive disorders	
Very rare:	Haemorrhagic corpus luteum / haemorrhagic ovarian cyst
Neoplasm benign, malignant and unspecified (including cysts and polyps)	
Rare:	Tumour lysis syndrome

1.Fatal cases have been reported in patients with advanced disease, severe infections, severe neutropenia and other serious concomitant conditions 2. Some fatal cases of gastrointestinal perforation have been reported

Description of selected adverse drug reactions

Myelosuppression

Myelosuppression is very common in cancer patients treated with Imatinib. Myelosuppression, thrombocytopenia, neutropenia and anaemia were the most frequently reported Grade 3 and 4 laboratory abnormalities. Overall, myelosuppression experienced with Imatinib in CML patients was generally reversible and in most patients did not result in dose interruption or dose reduction. Few patients required drug discontinuation. Other events of pancytopenia, lymphopenia and bone marrow depression have also been reported. Haematologic depression appeared greatest at the highest doses and also appeared to be dependent on the stage of CML disease, with Grade 3 or 4 neutropenia and thrombocytopenia between 4 and 6 times higher in blast and accelerated phase as compared to newly diagnosed patients in CP CML. These events can usually be managed with either a dose reduction or interruption, but they rarely require discontinuation of treatment with Imatinib. The incidence of hematologic toxicities is less in patients with solid tumours (i.e. GIST) than in patients with Ph+ leukaemias, with Grade 3/4 neutropenia and thrombocytopenia occurring approximately 10% and 1%, respectively.

Haemorrhage

CNS and GI haemorrhages are not uncommon in CML patients with compromised marrow function at baseline. Haemorrhages are well-recognized part of the disease complications in an acutely ill population of leukaemic patients, and may result from thrombocytopenia, or less commonly, platelet dysfunction. However, not all patients experiencing CNS and GI haemorrhages during therapy with Imatinib are thrombocytopenic. The most common manifestation of clinically significant bleeding was GI haemorrhage, which occurred most commonly in advanced CML patients and in metastatic GIST patients, where bleeding might occur as part of the underlying disease due to tumour bleeding from tumour haemorrhage/tumour necrosis. In first line CML and in adjuvant GIST setting, the observed frequencies of GI haemorrhage were generally the lowest. Gastric antral vascular ectasia (GAVE) is also rarely reported with Imatinib use in the post-marketing setting.

Oedema and Fluid Retention

Oedema is a common toxicity of Imatinib appearing in greater than 50% of all patients across all indications. Oedema is dose-related and there appears to be a correlation with its occurrence and plasma levels. The most common manifestation is periorbital oedema and somewhat less common is lower extremity oedema. Specific therapy is not usually required. Other fluid retention events occur much less commonly, but due to the location of the anatomic site may be potentially serious. The most frequent fluid retention event was pleural effusion, most commonly observed in advanced CML and metastatic GIST patients. The frequency of cardiac failure was generally low in patients with oedema and fluid retention.

It was higher in advanced CML than in other groups. This could be explained by the worse medical condition of advanced CML patients. The same trend was observed for renal failure in patients with oedema and fluid retention.

In a study, the frequency of events suggesting congestive heart failure was 1.5% on Imatinib vs. 1.1% on IFN-alpha in patients with newly-diagnosed CML. The frequency was appreciably higher in patients with transformed CML (accelerated phase or blast crisis), higher age, or with a baseline haemoglobin of less than 8 g/dL. Congestive Heart Failure (CHF) and left ventricular dysfunction have since been continuously monitored in the PSUR. Across all indications a higher frequency of CHF events observed in patients with CML than in patients with GIST might indicate differences of some of these disease-related risk factors. In addition, a published special safety analysis of cardiac events within the EORTC study of 942 patients with unresectable or metastatic GIST concluded that Imatinib does not induce left ventricular failure in GIST patients where the observed rate was approximately 0.2% while it can be up to 2% in a population with pre-existing cardiac disease.

Skin Rashes and Severe Cutaneous Adverse Reactions

A generalized erythematous, maculopapular, pruritic skin rash has been reported that can fade despite continued therapy. Some patients may have pruritus without accompanying rash, and sometimes there is an exfoliative component. Re-exposure in some patients has resulted in reappearance of rash, but not in all patients. These eruptions generally respond to antihistamines and topical steroids. Occasionally, systemic steroids are required. Skin rashes have been observed in up to one third of patients treated with imatinib across all indications. These are frequently pruritic and most commonly appear as erythematous, maculopapular or exfoliative lesions on the forearm, the trunk or the face or generalized with systemic expression. Skin biopsies have revealed a toxic drug reaction with a mixed cellular infiltrate. Although most rashes are mild and self-limiting more severe rare cases such as Stevens-Johnson toxic epidermal necrolysis, Erythema multiforme or DRESS may require interruption or discontinuation of treatment. Not surprisingly skin reactions were seen at a higher rate than placebo in the adjuvant GIST trial.

Hepatotoxicity

Hepatotoxicity, occasionally severe, may occur, and has been observed preclinically and clinically. LFT abnormalities usually consisted of mild elevations in transaminases, although a minority of patients had elevated levels of bilirubin. Onset is generally within the first two months of therapy, but has occurred as late as 6 to 12 months after commencing therapy. The levels generally normalize after withholding therapy for 1 to 4 weeks.

Hypophosphataemia

Low serum phosphate and hypophosphataemia (up to Grade 3 or 4) has been observed relatively commonly across all indications, however the origin and the clinical significance of this finding have not

been established. Imatinib has been shown to inhibit the differentiation of human monocytes into osteoclasts. The decrease was accompanied by a decrease in the resorptive capacity of these cells. A dose-dependent decrease of RANK-L was observed in osteoclasts in the presence of Imatinib. Sustained inhibition of osteoclastic activity may lead to counter regulatory response resulting in increased levels of PTH. The clinical relevance of the preclinical findings is yet unclear and an association with skeletal AEs such as bone fractures has not been demonstrated.

In the clinical development program serum phosphate was not routinely measured in all studies. Although it was initially hypothesized that hypophosphataemia might be dose-dependent, 24 month interpretable results from the Phase III TOPS study designed to investigate dose dependency of safety endpoints in patients with newly diagnosed CML, have shown that Grade 3 or 4 decreased serum phosphate or serum calcium has been experienced by 19.1% vs.15.5% and 5.1% vs. 0.9% of patients receiving 400 mg and 800 mg, respectively.

Gastrointestinal Obstruction, Perforation or Ulceration

GI ulceration, which may represent in extreme cases local irritation by Imatinib, has been observed in a small proportion of patients across all indications. Tumour haemorrhage/tumour necrosis, obstruction and GI perforation seem to be disease-related and have occurred exclusively or more frequently amongst GIST patients. In the case of metastatic GIST, tumor necrosis may occur in the context of tumor response, rarely leading to perforation. GI obstruction/ileus occurred most commonly in the GIST population where it may be caused by tumor obstruction from metastatic GIST and in the adjuvant setting by adhesions from previous GI surgery.

Tumour lysis syndrome

A causal relationship between tumour lysis syndrome and Imatinib treatment is deemed possible, although some cases were confounded by concomitant medications and other independent risks.

Growth retardation in pediatric patients

Imatinib appears to affect the stature of children, especially children who are pre-pubertal. A causal relationship between growth retardation in pediatric patients and Imatinib treatment could not be ruled out although for some cases of growth retardation in CML there was limited information.

Severe respiratory adverse drug reaction

Severe respiratory events, sometimes fatal, have been observed with Imatinib treatment, including acute respiratory failure, pulmonary hypertension, interstitial lung disease and pulmonary fibrosis. Pre-existing cardiac or pulmonary conditions that may be associated with severe respiratory events have been reported in many of these cases.

Laboratory test abnormalities

Haematology

CML-associated cytopenias, particularly neutropenia and thrombocytopenia, have been a consistent finding in all studies, with the suggestion of a higher frequency at high doses ≥750 mg (phase I study). However, the occurrence of cytopenias was also clearly dependent on the stage of the disease. In patients with newly diagnosed CML, cytopenias were less frequent than in the other CML patients. The frequency of Grade 3 or 4 neutropenias (ANC <1.0x10⁹/L) and thrombocytopenias (platelet count <50x10⁹/L) being between 4 and 6 times higher in blast crisis and accelerated phase (59 to 64% and 44 to 63% for neutropenia and thrombocytopenia, respectively) as compared to newly diagnosed patients in chronic phase CML (16.7% neutropenia and 8.9% thrombocytopenia). In newly diagnosed chronic phase CML Grade 4 neutropenia (ANC <0.5x10⁹/L) and thrombocytopenia (platelet count <10x10⁹/L) were observed in 3.6% and <1% of patients, respectively. The median duration of the neutropenic and thrombocytopenic episodes usually ranged from 2 to 3 weeks, and from 3 to 4 weeks, respectively. These events can usually be managed with either a dose reduction or an interruption of treatment with Imatinib, but can in rare cases lead to permanent discontinuation of treatment. In pediatric CML patients the most frequent toxicities observed were Grade 3 or 4 cytopenias involving neutropenia, thrombocytopenia and anaemia.

These generally occur within the first several months of therapy.

In patients with unresectable or metastatic malignant GIST, Grade 3 and 4 anaemias were reported in 5.4% and 0.7% of patients, respectively, and may have been related to gastrointestinal or intra-tumoural bleeding in at least some of these patients.

Grade 3 and 4 neutropenia were seen in 7.5% and 2.7% of patients, respectively, and Grade 3 thrombocytopenia in 0.7% of patients. No patient developed Grade 4 thrombocytopenia. The decreases in WBC and neutrophil counts occurred mainly during the first six weeks of therapy, with values remaining relatively stable thereafter.

Biochemistry

Severe elevation of transaminases (<5%) or bilirubin (<1%) has been seen in CML patients and was usually managed with dose reduction or interruption (the median duration of these episodes was approximately one week) of Imatinib. Treatment was discontinued permanently because of liver laboratory abnormalities in less than 1% of CML patients. In GIST patients, 6.8% of Grade 3 or 4 SGPT (serum glutamic pyruvic transferase) elevations and 4.8% of Grade 3 or 4 SGOT (serum glutamic oxaloacetic transferase) elevations were observed. Bilirubin elevation was below 3%.

There have been cases of cytolytic and cholestatic hepatitis and hepatic failure; in some of which outcome was fatal.

4.9 Overdose

Experience with higher than therapeutic doses is limited. Isolated cases of Imatinib overdose have been reported spontaneously and in the literature. Generally the reported outcome in these cases was improvement or recovery. In the event of overdose the patient should be observed and appropriate symptomatic treatment should be given.

Events that have been reported at different dose ranges are as follows:

Adult overdose

1,200 to 1,600 mg (duration varying between 1 to 10 days): Nausea, vomiting, diarrhea, rash, erythema, edema, swelling, fatigue, muscle spasms, thrombocytopenia, pancytopenia, abdominal pain, headache, decreased appetite.

1,800 to 3,200 mg (as high as 3,200 mg daily for 6 days): Weakness, myalgia, increased CPK, increased bilirubin, gastrointestinal pain.

6,400 mg (single dose): One case in the literature reported one patient who experienced nausea, vomiting, abdominal pain, pyrexia, facial swelling, neutrophil count decreased, increased transaminases.

8 to 10 g (single dose): Vomiting and gastrointestinal pain have been reported.

Pediatric overdose

One 3 year-old male exposed to a single dose of 400 mg experienced vomiting, diarrhea and anorexia and another 3 year old male exposed to a single dose of 980 mg dose experienced decreased white blood cell count and diarrhea.

5. PHARMACOLOGICAL PROPERTIES

Pharmacotherapeutic group: Antineoplastic agents, protein kinase inhibitors, ATC code: L01EA01."

5.1 Pharmacodynamic Properties

Mechanism of action (MOA)

Imatinib is a small molecule protein-tyrosine kinase inhibitor that potently inhibits the activity of the BCR-ABL tyrosine kinase (TK), as well as several receptor TKs: KIT, the receptor for stem cell factor (SCF) coded for by the KIT proto-oncogene, the discoidin domain receptors (DDR1) and DDR2), the colony stimulating factor receptor (CSF-1R) and the platelet-derived growth factor receptors alpha and beta (PDGFR-alpha and PDGFR-beta). Imatinib can also inhibit cellular events mediated by activation of these receptor kinases.

Pharmacodynamics (PD)

Imatinib is a protein-tyrosine kinase inhibitor, which potently inhibits the breakpoint cluster

region-Abelson (BCR-ABL) tyrosine kinase at the in vitro, cellular, in vivo levels. The compound selectively inhibits proliferation and induces apoptosis in BCR-ABL positive cell lines as well as fresh leukemic cells from Philadelphia chromosome positive CML and acute lymphoblastic leukemia (ALL) patients. In colony transformation assays using ex vivo peripheral blood and bone marrow samples, Imatinib shows selective inhibition of BCR-ABL positive colonies from CML patients.

In vivo the compound shows anti-tumor activity as a single agent in animal models using BCR-ABL positive tumor cells. Imatinib is also an inhibitor of the receptor tyrosine kinases for platelet-derived growth factor (PDGF) and stem cell factor (SCF), KIT, and inhibits PDGF- and SCF-mediated cellular events. In vitro, Imatinib inhibits proliferation and induces apoptosis in gastrointestinal stromal tumor (GIST) cells, which express an activating KIT mutation. Constitutive activation of the PDGFR or the ABL protein tyrosine kinases as a consequence of fusion to diverse partner proteins or constitutive production of PDGF have been implicated in the pathogenesis of MDS/MPD, HES/CEL and DFSP. In addition, constitutive activation of KIT or the PDGFR has been implicated in the pathogenesis of ASM. Imatinib inhibits signaling and proliferation of cells driven by dysregulated PDGFR, KIT and ABL kinase activity.

5.2 Pharmacokinetic Properties

The pharmacokinetics of Imatinib have been evaluated over a dosage range of 25 to 1,000 mg. Plasma pharmacokinetic profiles were analyzed on day 1 and on either day 7 or day 28, by which time plasma concentrations had reached steady state.

Absorption

Mean absolute bioavailability for the tablet formulation imatinib is 98%. The coefficient of variation for plasma imatinib AUC is in the range of 40% to 60% after an oral dose. When given with a high fat meal, the rate of absorption of Imatinib was minimally reduced (11% decrease in Cmax and prolongation of tmax by 1.5 h), with a small reduction in AUC (7.4%) compared to fasting conditions.

Distribution

At clinically relevant concentrations of Imatinib, binding to plasma proteins was approximately 95% on the basis of in vitro experiments, mostly to albumin and alpha-acid-glycoprotein, with little binding to lipoprotein.

Biotransformation/metabolism

The main circulating metabolite in humans is the N-demethylated piperazine derivative (CGP71588), which shows similar in vitro potency as the parent compound. The plasma AUC for this metabolite was found to be only 16% of the AUC for Imatinib. The plasma protein binding of the N-demethylated metabolite is similar to that of the parent compound.

Elimination

Based on the recovery of compound(s) after an oral ¹⁴C-labelled dose of Imatinib, approximately 81% of the dose was eliminated within 7 days in feces (68% of dose) and urine (13% of dose). Unchanged Imatinib accounted for 25% of the dose (5% urine, 20% feces), the remainder being metabolites. The mean apparent elimination half-life estimated from the single dose PK study was 13.5 hours. The half-life of all ¹⁴C-labelled components in plasma was from 41-72 hours.

Plasma pharmacokinetics

Following oral administration in healthy volunteers, the t^{1/2} was approximately 18 h, suggesting that once-daily dosing is appropriate. The increase in mean AUC with increasing dose was linear and dose proportional in the range of 25 to 1,000 mg Imatinib after oral administration. There was no change in the kinetics of imatinib on repeated dosing, and accumulation was 1.5 to 2.5 fold at steady state when dosed once daily.

Special populations

Based on population pharmacokinetic analysis, there was a small effect of age on the volume of distribution (12% increase in patients >65 years old). This change is not thought to be clinically significant. The effect of body weight on the clearance of Imatinib is such that for a patient weighing 50 kg the mean clearance is expected to be 8.5 L/h, while for a patient weighing 100 kg the clearance will rise to 11.8 L/h. These changes are not considered sufficient to warrant dose adjustment based on kg bodyweight. There is no effect of gender on the kinetics of Imatinib.

Further population PK analysis in the study in newly diagnosed CML patients showed that the effect of covariates and co-medications on both clearance and volume of distribution appears to be small and is not sufficiently pronounced to warrant dose adjustment.

Pediatric patients (below 18 years)

As in adult patients, Imatinib was rapidly absorbed after oral administration in pediatric patients in studies. Dosing in children at 260 and 340 mg/m² achieved the same exposure, respectively, as doses of 400 mg and 600 mg in adult patients. The comparison of AUC⁰⁻²⁴ on Day 8 and Day 1 at 340 mg/m² dose level revealed a 1.7 fold drug accumulation after repeated once daily dosing.

Based on pooled population pharmacokinetic analysis in pediatric patients with hematological disorders (CML, Ph+ALL, or other hematological disorders treated with Imatinib), clearance of Imatinib increases with increasing body surface area (BSA). After correcting for the BSA effect, other demographics such as age, body weight and body mass index did not have clinically significant effects on the exposure of Imatinib. The analysis confirmed that exposure of Imatinib in pediatric patients receiving 260 mg/m² once daily (not exceeding 400 mg once daily) or 340 mg/m² once daily (not exceeding 600 mg once daily) were similar to those in adult patients who received imatinib 400 mg or 600 mg once daily.

Organ function impairment

Imatinib and its metabolites are not excreted via the kidney to a significant extent. Patients with mild and moderate impairment of renal function appear to have a higher plasma exposure than patients with normal renal function. The increase is approximately 1.5 to 2 fold, corresponding to a 1.5 fold elevation of plasma AGP, to which Imatinib binds strongly. The free drug clearance of Imatinib is probably similar between patients with renal impairment and those with normal renal function, since renal excretion represents only a minor elimination pathway for Imatinib.

Although the results of pharmacokinetic analysis showed that there is considerable inter-subject variation, the mean exposure to Imatinib did not increase in patients with varying degrees of liver dysfunction as compared to patients with normal liver function.

6. PHARMACEUTICAL PARTICULARS

6.1 Pack Size

For 100mg tablets: 6 x 10's (60 tablets)

For 400mg tablets: 3 x 10's (30 tablets)

Alu-Alu blisters having 10 tablets in each blister and packed in unit carton with leaflet.

6.2 Storage Conditions

Store below 30°C. Protect from sunlight and moisture.

6.3 Dosage and Instructions

As directed by physician.

Keep out of reach of children.

For adverse event reporting, kindly email us at

pharmacovigilance@oncogenpharma.com.pk

7. MANUFACTURER

Manufactured by

oncogen

Oncogen Pharma (Pvt.) Limited,

Plot No: WII-26 & 27-A, Korangi Creek Industrial Park (KCIP), Karachi 75190, Pakistan,

www.oncogenpharma.com.pk

خوداک: ذاکر کی ہدایت کے مطابق استعمال کریں۔

ہدایت: دو اکو سو ڈگری سینٹی گریڈ سے کم درجہ حرارت پر رکھیں۔

دعویٰ اور نی سے بچائیں۔

بچوں کی پہنچ سے دور رکھیں۔